

PARTON SATURATION-AN OVERVIEW¹

A.H. Mueller²

Department of Physics, Columbia University
New York, New York 10027

The idea of partons and the utility of using light-cone gauge in QCD are introduced. Saturation of quark and gluon distributions are discussed using simple models and in a more general context. The Golec-Biernat Wüsthoff model and some simple phenomenology are described. A simple, but realistic, equation for unitarity, the Kovchegov equation, is discussed, and an elementary derivation of the JIMWLK equation is given.

1 Introduction

These lectures are meant to be an introduction, and an overview, of parton saturation in QCD. Parton saturation is the idea that the occupation numbers of small- x quarks and gluons cannot become arbitrarily large in the light-cone wavefunction of a hadron or nucleus. Parton saturation is an idea which is becoming well established theoretically and has important applications in small- x physics in high-energy lepton-hadron collisions and in the early stages of high-energy heavy ion collisions. The current experimental situation is unclear. Although saturation based models have had considerable phenomenological success in explaining data at HERA and at RHIC more complete and decisive tests are necessary before it can be concluded that parton saturation has been seen. These lectures begin very simply by describing, through an example, some important features of light-cone perturbation theory in QCD, and they end by describing some rather sophisticated equations which govern light-cone wavefunctions when parton densities are very large.

¹Lectures given at the Cargese Summer School, August 6-18, 2001

²This work is supported in part by the Department of Energy

$$|\Psi_p\rangle = N \left(\text{quark } p \right) + \text{quark } p \text{ emitting gluon } (k, c, \lambda) \text{ and continuing as } (p-k)$$

FIG.1

2 States in QCD Perturbation Theory

In light-cone perturbation theory states of QCD are described in terms of the numbers and distributions in momentum of quarks and gluons. The essential features of light-cone perturbation theory that will be needed in these lectures can be illustrated by considering the wavefunction of a quark through lowest order in the QCD coupling g . One can write

$$|\psi_p\rangle = N|p\rangle + \sum_{\lambda=\pm} \sum_{c=1}^{N_c^2-1} \int d^3k \psi_\lambda^c(k) |p-k; k(\lambda, c)\rangle \quad (1)$$

where $|p\rangle$ is a free quark state with momentum p , $|\psi_p\rangle$ is a dressed quark state, and $|p-k; k(\lambda, c)\rangle$ is a state of a quark, of momentum $p-k$, and a gluon of momentum k , helicity λ and color c . We suppress quark color indices which will appear in matrix form in what follows. We label states by momenta $p_+ = \frac{1}{\sqrt{2}}(p_0 + p_3)$, p_1, p_2 , and $d^3k = dk_+ d^2k = dk_+ dk_1 dk_2$. Recall that in light-cone quantization momenta P_+ and $\underline{P} = (P_1, P_2)$ are kinematic while P_- plays the role of a Hamiltonian and generates evolution in the “time” variable $x_+ = \frac{1}{\sqrt{2}}(x_0 + x_3)$. For an on-shell zero mass particle $p_- = \underline{p}^2/2p_+$. N in (1) is a normalization factor. Eq.(1) is illustrated in Fig.1.

ψ_λ^c is determined from light-cone perturbation theory to be

$$\delta^3(p-p')\psi_\lambda^c(k) = \frac{\langle p'-k; k(\lambda, c) | H_I | p \rangle}{(p'-k)_- + k_- - p_-} \quad (2)$$

with

$$H_I = g \int d^3x \bar{q}(x) \gamma_\mu \left(\frac{\lambda^c}{2} \right) q(x) A_\mu^c(x) \quad (3)$$

where the gluon field is

$$A_\mu^c(x) = \sum_{\lambda=\pm} \int \frac{d^3k}{\sqrt{(2\pi)^3 2k_+}} [\epsilon_\mu^\lambda(k) a_\lambda^c(k) e^{i\mathbf{k}\cdot\mathbf{x} - ik_+x_- - ik_-x_+} + h.c.] \quad (4)$$

with

$$[a_{\lambda'}^{c'}(k'), a_\lambda^{\dagger c}(k)] = \delta_{\lambda'\lambda} \delta_{c'c} \delta^3(k' - k). \quad (5)$$

In (2) and (5) $\delta^3(p) = \delta(p_+) \delta^2(\underline{p})$ while $d^3x = dx_- d^2\underline{x}$ in (3). It is useful to imagine the calculation being done in a frame where p_+ is large and $\underline{p} = 0$ in which case

$$k_- = \frac{\underline{k}^2}{2k_+}, \quad (p-k)_- = \frac{\underline{k}^2}{2(p-k)_+}. \quad (6)$$

In the soft gluon approximation $k_+/p_+ \ll 1$ and thus $k_- \gg (p-k)_-$ so that only k_- need be kept in the denominator in (2). In addition, in light-cone gauge, $A_+ = 0$, the polarization vectors can be written as

$$\epsilon_\mu^\lambda(k) = (\epsilon_+^\lambda, \epsilon_-^\lambda, \underline{\epsilon}^\lambda) = (0, \frac{\underline{\epsilon}^\lambda \cdot \underline{k}}{k_+}, \underline{\epsilon}^\lambda) \quad (7)$$

and, because of the $1/k_+$ term, only ϵ_-^λ need be kept in (2) in the soft gluon approximation. Using (3)-(7) in (2) one finds

$$\psi_\lambda^c(k) = \left(\frac{\lambda^c}{2}\right) 2g \frac{(\underline{\epsilon}^\lambda)^* \cdot \underline{k}}{\underline{k}^2} \frac{1}{\sqrt{(2\pi)^3 2k_+}} \quad (8)$$

Problem 1(E): Using the formula $\bar{U}(p-k)\gamma_\mu\bar{U}(p) \approx 2p_+g_{\mu-}$ for high momentum Dirac spinors derive (8).

3 Partons

Define the gluon distribution of a state $|S(p)\rangle$ by

$$xG_S(x, Q^2) = \sum_{\lambda, c} \int d^3k x \delta(x - k_+/p_+) \Theta(Q^2 - \underline{k}^2) \langle S(p) | a_\lambda^{c\dagger}(k) a_\lambda^c(k) | S(p) \rangle. \quad (9)$$

The meaning of xG_S is clear. $xG_S(x, Q^2)dx$ is the number of gluons, having longitudinal momentum between xp_+ and $(x + dx)p_+$, localized in transverse coordinate space to a region $\Delta x_\perp \sim 1/Q$, in the state $|S(p)\rangle$. For a quark, at order g^2 , one finds from (1)

$$xG_q(x, Q^2) = \sum_{\lambda, c} \int d^3k x \delta(x - k_+/p_+) \Theta(Q^2 - \underline{k}^2) \psi_\lambda^{c\dagger}(k) \psi_\lambda^c(k). \quad (10)$$

Using (8) in (10) one finds

$$xG_q(x, Q^2) = \sum_c \frac{\lambda^c}{2} \frac{\lambda^c}{2} \int \frac{4g^2}{(2\pi)^3} \frac{d^2k}{\underline{k}^2} \frac{dk_+}{2k_+} x \delta(x - k_+/p_+) \Theta(Q^2 - \underline{k}^2). \quad (11)$$

Using $\sum_c \frac{\lambda^c}{2} \frac{\lambda^c}{2} = C_F = \frac{N_c^2 - 1}{2N_c}$ and introducing an infrared cutoff, μ , for the transverse momentum integral in (11) gives

$$xG_q(x, Q^2) = \frac{\alpha C_F}{\pi} \ell n(Q^2/\mu^2). \quad (12)$$

We note that if $xG(x, Q^2) = 3xG_q(x, Q^2)$ is taken one obtains a result for the proton which is not unreasonable phenomenologically for $x \sim 10^{-2} - 10^{-1}$ and moderate Q^2 if μ is taken to be 100 MeV.

4 Classical Fields

One can associate a classical field with gluons in the quark.

$$A_i^{(c\ell)}(x) = \int d^3p' \langle \psi_{p'} | A_i^c(x) | \psi_p \rangle. \quad (13)$$

Using (1) and (8) in (13) one finds

$$A_i^{(c\ell)}(x) = \int \frac{d^3k}{(2\pi)^3} e^{-ik \cdot x} \left(\frac{\lambda^c}{2} \right) \frac{g k_i}{\underline{k}^2 k_+} \quad (14)$$

or

$$A_i^{c(\ell)}(x) = \int \frac{d^3k}{(2\pi)^3} e^{-ik \cdot x} A_i^{c(\ell)}(k) \quad (15)$$

with

$$A_i^{c(\ell)}(k) = \frac{\lambda^c}{2} \frac{gk_i}{k_+}. \quad (16)$$

In (14) the k_+ integration goes from $-\infty$ to $+\infty$. The region $k_+ > 0$ comes from $|\psi_p\rangle$ consisting of a bare quark and a gluon of momentum k while the region $k_+ < 0$ comes from $|\psi_{p'}\rangle$ consisting of a bare quark and a gluon.

Problem 2(E): Take $\frac{1}{k_+} = \frac{1}{k_+ - i\epsilon}$ in (16) and show that

$$A_i^{c(\ell)}(x) = -g\left(\frac{\lambda^2}{2}\right) \frac{x_i}{2\pi \underline{x}^2} \Theta(-x_-) \text{ and } F_{+i}^{c(\ell)} = \frac{\partial}{\partial x_-} A_i^{c(\ell)} = g\left(\frac{\lambda^c}{2}\right) \frac{x_i}{2\pi \underline{x}^2} \delta(x_-).$$

Problem 3(E): Show that G_q , as given in (10), can also be written as

$$xG_q(x, Q^2) = \int \frac{d^3k}{(2\pi)^3} \delta(x_- k_+ / p_+) \Theta(Q^2 - \underline{k}^2) \sum_{i,c} [A_i^{c(\ell)}(k)]^2 2k_+.$$

5 Why Light-Cone Gauge is Special

In order to understand why light-cone gauge plays a special role in describing high-energy hadronic states a simple calculation of the near forward high-energy elastic amplitude for electron-electron scattering in QED is useful. The graph to be calculated is shown in Fig.2 and we imagine the calculation being done in the center of mass frame with p being a right-mover (p_+ large) and p_1 being a left-mover (p_{1-} large). First we shall do the calculation in covariant gauge and afterwards in light-cone gauge.

In covariant gauge the photon propagator, the k -line in Fig.2, is $D_{\mu\nu} = \frac{-i}{k^2} g_{\mu\nu}$ and the dominant term comes from taking $g_{\mu\nu} \rightarrow g_{-+} = 1$ giving

$$T = ig \frac{\bar{U}(p-k)\gamma_+ U(p)}{2p_+} ig \frac{\bar{U}(p_1+k)\gamma_- U(p_1)}{2p_{1-}} \frac{-i}{k^2}. \quad (17)$$




FIG.2

Using $\bar{U}(p-k)\gamma_+U(p) \approx 2p_+$ and $\bar{U}(p_1+k)\gamma_-U(p_1) \approx 2p_{1-}$ one finds

$$T = \frac{-ig^2}{\underline{k}^2} \quad (18)$$

where we have used $k^2 = 2k_+k_- - \underline{k}^2 \approx -\underline{k}^2$ since both k_+ and k_- are required to be small from the mass shell conditions $(p-k)^2 = (p_1+k)^2 = 0$ for the zero mass electrons.

Now suppose we do the calculation in light-cone gauge $A_+ = 0$ where the propagator is

$$D_{\mu\nu}(k) = \frac{-i}{k^2} [g_{\mu\nu} - \frac{\eta_\mu k_\nu + \eta_\nu k_\mu}{\eta \cdot k}] \quad (19)$$

where $\eta \cdot V = V_+$ for any vector V_μ . Now the dominant term comes from taking D_{-i} giving

$$T = iq \frac{\bar{U}(p-k)\gamma_+U(p)}{2p_+} ig \frac{\bar{U}(p_1+k)\underline{\gamma} \cdot \underline{k} U(p_1)}{2p_{1-}} \frac{-i}{k_+k^2}. \quad (20)$$

Using $\bar{U}(p_1+k)\underline{\gamma} \cdot \underline{k} U(p_1) = \underline{k}^2$ one finds

$$T = \frac{-ig^2}{2p_{1-}k_+}. \quad (21)$$

Now $(p_1+k)^2 = 0$ gives $2p_{1-}k_+ \approx \underline{k}^2$ so that (18) and (21) agree as expected.

The result (18) comes about in a natural way $\frac{\bar{U}(p-k)\gamma_+U(p)}{2p_+}$ and $\frac{\bar{U}(p_1+k)\gamma_-U(p_1)}{2p_{1-}}$ are the classical currents, equal to 1, of particles moving along the light-cone while the $1/\underline{k}^2$ factor is just the (instantaneous) potential between the

charges. However, when one uses $A_+ = 0$ light-cone gauge the dominant part of the current for left moving particles is forbidden and one must keep the small transverse current $\underline{k}/2p_{1-}$. However, the smallness of the current is compensated by the factor $1/k_+$ in the light-cone gauge propagator. In coordinate space the $1/k_+$ comes about from the potential acting over distances $x_- \approx 1/k_+ = \frac{2p_{1-}}{k^2}$ so that the potential is very non-local and non-causal. By choice of the $i\epsilon$ prescription[1, 2] for $1/k_+$ one can put these non-causal interactions completely before the scattering (initial state) or completely after the scattering (final state). In Problem 2 the potential A_i exhibits this non-causal behavior with our choice of $i\epsilon$ placing the long time behavior in the initial state, the $\Theta(-x_-)$ term.

6 High Momentum Particles and Wilson Lines³⁻⁵

Suppose a quark of momentum p is a high-energy right mover, that is $p_+ \gg p_-, \underline{p}$. Then so long as one does not choose to work in $A_- = 0$ light-cone gauge the dominant coupling of gauge fields to p are classical (eikonal) when p passes some QCD hadron or source. To be specific suppose p scatters on a hadron elastically and with a small momentum transfer. We may view the interactions as shown in Fig.3 for a three-gluon exchange term. Of course to get the complete scattering one has to sum over all numbers of gluon exchanges. Call $S(p_+, \underline{b})$ the S-matrix for scattering of the right moving quark on the target. (We imagine that the target hadron has large gluon fields making it necessary to find a formula which includes all gluon interactions.) Although the right moving quark has sufficient momentum so that it moves close to the light-cone we do assume that the momentum is not so large that higher gluonic components of the quark wavefunction need be considered. Since the probability of extra transverse gluons being present in the wavefunction is in general proportional to αy , with y being the quark rapidity, we suppose $\alpha y \ll 1$.

The graph shown in Fig.3 can be written as




FIG.3.

$$S(p_+, \underline{b}) = \int \left\{ \frac{1}{\sqrt{2(p+k)_+}} \bar{U}(p+k) i g T^a \gamma_+ \frac{i}{\gamma \cdot (p+k_1+k_2)} i g T^b \gamma_+ \frac{i}{\gamma \cdot (p+k_1)} \right. \\ \left. \cdot i g T^c \gamma_+ U(p) \frac{1}{\sqrt{(2p_+)}} \right\} \frac{d^2 k d k_+}{(2\pi)^3} e^{i \underline{k} \cdot \underline{b}} \frac{d^4 k_1 d^4 k_2}{(2\pi)^8} M_{abc}(k_1, k_2) \quad (22)$$

where

$$M_{abc}(k_1, k_2) = \int d^4 x_1 d^4 x_2 d^4 x e^{i(k-k_1-k_2) \cdot x + i k_2 \cdot x_2 + i k_1 \cdot x_1} \\ \cdot \langle f | T A_+^a(x) A_+^b(x_2) A_+^c(x_1) | i \rangle. \quad (23)$$

It is straightforward to evaluate the $\{ \}$ -term in (22) and one gets

$$\{ \} = i g T^a i g T^b i g T^c \frac{i}{k_{1-} + i\epsilon} \frac{i}{(k_1 + k_2)_- + i\epsilon}. \quad (24)$$

Now do the $d^2 k_1 d k_{1+}$ and $d^2 k_2 d k_{2+}$ integrals followed by $d^2 x_1 d x_{1-}$ and $d^2 x_2 d x_{2-}$. This sets $\underline{x}_1 = \underline{x}_2 = \underline{x}$ and $x_{1-} = x_{2-} = x_-$. The only non-zero term in the time-ordered product is proportional to $\Theta(x_- - x_{2-}) \Theta(x_{2-} - x_{1-})$ and this factor along with the exponentials in x_{1-}, x_{2-} and x_- allow the $d k_{1-}$ and $d k_{2-}$ integrals to be done over the poles in (24). We get finally

$$S_{fi}(p_+, \underline{b}) = (ig)^3 \langle f | \int_{-\infty}^{\infty} d x_+ A_-(\underline{b}, x_+) \int_{-\infty}^{x_+} d x_{1+} A_-(\underline{b}, x_{2+}) \\ \cdot \int_{-\infty}^{x_{2+}} d x_{1+} A_-(\underline{b}, x_{1+}) | i \rangle \quad (25)$$

where $A_\mu = T^a A_\mu^a$ and we have suppressed the variable $x_- = 0$ in the A' s in (25).

The general term is now apparent, and one has in the general case

$$S_{fi}(p_+, \underline{b}) = \langle f | P e^{ig \int_{-\infty}^{\infty} dx_+ A_-(\underline{b}, x_+)} | i \rangle \quad (26)$$

where P denote an x_+ - ordering of the matrices A where A' s having larger values of x_+ come to the left of those having smaller values.

Problem 4(M): Show that the time orderings different from $\Theta(x_+ - x_{2+})\Theta(x_{2+} - x_{1+})$ give no contribution to (22).

Finally, a word of caution in using (26). In general there are singularities present when two values of x_+ , in adjoining A' s, become equal, although in many simple models the x_+ - integrations are regular. If singularities in the x_+ integrations arise it is generally possible to extract the leading logarithmic contributions to the scattering amplitude by carefully examining the singularities[5]. A detailed discussion of this is, however, far beyond the scope of these lectures.

7 Dual Descriptions of Deep Inelastic Scattering; Bjorken and Dipole Frames

Particular insight into the dynamics of a process often occurs by choosing a particular frame and an appropriate gauge. Indeed the physical picture of a process may change dramatically in different frames and in different gauges. In a frame where the parton picture of a hadron is manifest saturation shows up as a limit on the occupation number of quarks and gluons, however, in a different (dual) frame saturation appears as the unitarity limit for scattering of a quark or of a gluon dipole at high-energy[6].

To see all this a bit more clearly consider inelastic lepton-nucleon scattering as illustrated in Fig.4 where a lepton emits a virtual photon which then scatters on a nucleon. We suppose $Q^2 = -q^2$ is large. The two structure functions, F_1 and F_2 , which describe the cross section can depend on the invariants Q^2 and $x = \frac{Q^2}{2P \cdot q}$. To make the parton picture manifest we choose $A_+ = 0$ light-cone gauge along with the frame




FIG.4

$$P = (P + \frac{M^2}{2P}, 0, 0, P)$$

and

$$q = (q_0, \underline{q}, q_z = 0),$$

and where $P \rightarrow \infty$. This is the Bjorken frame. We note that $q_0 = \frac{P \cdot q}{P}$ goes to zero as $P \rightarrow \infty$ so that the virtual photon momentum is mainly transverse to the nucleon direction. This last fact means that the virtual photon is a good analyzer of transverse structure since it is absorbed over a transverse distance $\Delta x_\perp \sim 1/Q$. Since $\Delta x_\perp$ is very small at large Q the virtual photon is absorbed by, and measures, individual quarks.

Problem 5(H): Use the uncertainty principle to show that the time, $\Delta\tau$, over which $\gamma^*(q)$ is absorbed by a quark is $\Delta\tau \approx 2xP/Q^2$. You may assume that $\underline{k}^2/Q^2 \ll 1$ where the process of absorption of the photon is illustrated in Fig.5.

The result of problem 5 and of the result that $\Delta x_\perp \sim 1/Q$ motivates the formula




FIG.5

$$F_2(x, Q^2) = \sum_f e_f^2 [xq_f(x, Q^2) + x\bar{q}_f(x, Q^2)] \quad (27)$$

which says that the structure function F_2 is proportional to the charge squared of the quark, having flavor f , absorbing the photon and proportional to the number of quarks having longitudinal momentum fraction x and localized in transverse coordinate space to a size $1/Q$. F_2 is given in terms of the longitudinal and transverse virtual photon cross sections on the proton as

$$F_2 = \frac{1}{4\pi^2\alpha_{em}} Q^2 [\sigma_T + \sigma_L]. \quad (28)$$

Eq.(27) is the QCD improved parton model. In more technical terms Q^2 is a renormalization point which in the parton picture is a cutoff of the type given in (9) for gluon distributions and here occurring for quark distributions.

Now consider the same process in the dipole frame pictured in Fig.6 where

$$P = (P + \frac{M^2}{2P}, 0, 0, P)$$

$$q = (\sqrt{q^2 - Q^2}, 0, 0, -q)$$




FIG.6

and where $q/Q \gg 1$ but where q is fixed as x becomes small so that most of the energy in a small- x process, and it is only for small- x scattering that the dipole frame is useful, still is carried by the proton. Now we suppose that a gauge different from $A_+ = 0$ is being used, for example a covariant gauge or the gauge $A_- = 0$. In the dipole frame the process looks like $\gamma^* \rightarrow$ quark – antiquark followed by the scattering of the quark-antiquark dipole on the nucleon[7, 8, 9]. The splitting of γ^* into the quark-antiquark pair is given by lowest order perturbation theory while all the dynamics is in the dipole-nucleon scattering. In this frame the partonic structure of the nucleon is no longer manifest, and the virtual photon no longer acts as a probe of the nucleon.

Equating these two pictures and fixing the transverse momentum of the leading quark or antiquark one has[6]

$$e_f^2 \frac{d(xq_f + x\bar{q}_f)}{d^2b d^2\ell} = \frac{Q^2}{4\pi^2 \alpha_{em}} \int \frac{d^2x_1 d^2x_2}{4\pi^2} \int_0^1 dz \frac{1}{2} \sum_{\lambda} \psi_{T\lambda}^{f*}(\underline{x}_2, z, Q) \psi_{T\lambda}^f(\underline{x}_1, z, Q) \cdot e^{-i\ell \cdot (\underline{x}_1 - \underline{x}_2)} [S^\dagger(\underline{x}_2) S(\underline{x}_1) - S^\dagger(\underline{x}_2) - S(\underline{x}_1) + 1], \quad (29)$$

where the γ^* wavefunction is

$$\psi_{T\lambda}^f(\underline{x}, z, Q) = \left\{ \frac{\alpha_{em} N_c}{2\pi^2} z(1-z)[z^2 + (1-z)^2] Q^2 \right\}^{1/2} e_f K_1(\sqrt{Q^2 \underline{x}^2 z(1-z)}) \frac{\underline{\epsilon}^\lambda \cdot \underline{x}}{|\underline{x}|}. \quad (30)$$

We have written (29) for a fixed impact parameter, $\underline{b}$, which is a (suppressed) variable in S . To make the identification exhibited in (29) requires that the struck quark shown in Fig.5 not have final state interactions, and this requires the special choice of $i\epsilon$'s in the light-cone gauge, $A_+ = 0$, used to identify the quark distributions. With this choice of $i\epsilon$'s the lefthand side of (29) refers to the density of quarks in the nucleon wavefunction while the righthand side of (29) refers to the transverse momentum spectrum of quark jets produced. This identification is only possible when final state interactions are absent. Finally, it is not hard to see the origin of the final factor on the righthand side of (29). This is just $[S^\dagger(\underline{x}_2) - 1][S(\underline{x}_1) - 1]$, the product of the $T-$ matrices for the dipole $\underline{x}_1$ in the amplitude and the dipole $\underline{x}_2$ in the complex conjugate amplitude.

8 Quark Distributions in a large Nucleus; Quark Saturation at One-Loop

In general it is very difficult to evaluate the $S-$ matrices appearing in (29). However, if the target is a large nucleus one can define an interesting model, if not a realistic calculation for real nuclei, by limiting the dipole nucleon interaction to one and two gluon exchanges. That is, we suppose the dipole-nucleon interaction is weak. Despite this assumed weakness of interaction the dipole nucleus scattering can be very strong if the nucleus is large enough. We begin with the term $S(\underline{x}_1)$ in (29), an elastic scattering of the dipole on the nucleus in the amplitude with no interaction at all in the complex conjugate amplitude. This term is illustrated in Fig.7.

Now $|S(\underline{x}_1)|^2$ is the probability that the dipole does not have an inelastic interaction as it passes through the nucleus. We can write

$$|S(\underline{x}_1)|^2 = e^{-L/\lambda} \quad (31)$$

where $L = 2\sqrt{R^2 - \underline{b}^2}$ is the length of nuclear matter that the dipole traverses at impact parameter $\underline{b}$ for a uniform spherical nucleus of radius R , and λ is the mean free path for inelastic dipole-nucleon interactions. Using




FIG.7

$$\lambda = [\rho\sigma]^{-1} \quad (32)$$

with ρ the nuclear density one has

$$S(\underline{x}_1, \underline{b}) = e^{-2\sqrt{R^2 - b^2}\rho\sigma(\underline{x}_1)/2} \quad (33)$$

if we suppose S is purely real. Detailed calculation gives[10]

$$\sigma(\underline{x}) = \frac{\pi^2\alpha}{N_c} xG(x, 1/\underline{x}^2)\underline{x}^2. \quad (34)$$

Problem 6(H*): Check (34) for scattering of a dipole on a bare quark.

Thus we know how to calculate the last three terms in [] in (29). What about the $S^\dagger S$ term? Graphically this term is illustrated in Fig.8 for some typical elastic and inelastic interactions of the dipoles with the nucleons in the nucleus. Let us focus on the final interaction, the one closest to the cut (the vertical line) in Fig.8. One can check that interactions with the $\underline{x} = \underline{0}$ line cancel between interactions in the amplitude and the complex conjugate amplitude.

Problem 7(H): Verify, using $S^\dagger S = 1$, that the three interactions shown in Fig.9 cancel. You may assume that T is purely imaginary ($S = 1 - iT$) although this is not necessary for the result.




FIG.8




FIG.9




FIG.10




FIG.11a




FIG.11b.

Problem 8(H): Again, using $S^\dagger S = 1$, show that the two interactions shown in Fig.10 cancel. Assume T is purely imaginary.

The results of problems 7 and 8 establish that one need not consider interactions with the $\underline{x} = 0$ line in evaluating the $S^\dagger(\underline{x}_2)S(\underline{x}_1)$ term in (29). Thus, we are left only with interactions on the $\underline{x}_1$ -line in the amplitude and with the $\underline{x}_2$ -line in the complex conjugate amplitude. The result of problem 8 allows one to transfer the interactions with the $\underline{x}_2$ -line in the complex conjugate amplitude to interactions with a line placed at $\underline{x}_2$ in the amplitude[11], although this does require that S be real. This is illustrated in the equality between the terms in Fig.11a and Fig.11b and reads

$$S^\dagger(\underline{x}_2)S(\underline{x}_1) = S(\underline{x}_1 - \underline{x}_2). \quad (35)$$

Eq.(35) is a beautiful result, but unfortunately it likely has a limited validity. It appears to require a maximum of two gluon lines exchanged between any given nucleon and the dipole lines as well as the requirement of a purely real S-matrix for elastic scattering. Thus using (29), (33) and (35) we arrive at[6]

$$e_f^2 \frac{d(xq_f + x\bar{q}_f)_A}{d^2\ell d^2b} = \frac{Q^2}{4\pi^2\alpha_{em}} \int \frac{d^2x_1 d^2x_2}{4\pi^2} \int_0^1 dz \frac{1}{2} \sum_\lambda \psi_{T\lambda}^{f*} \psi_{T\lambda}^f e^{-i\ell \cdot (\underline{x}_1 - \underline{x}_2)} \cdot [1 + e^{-(\underline{x}_1 - \underline{x}_2)^2 \bar{Q}_S^2/4} - e^{-\underline{x}_1^2 \bar{Q}_S^2/4} - e^{-\underline{x}_2^2 \bar{Q}_S^2/4}] \quad (36)$$

with

$$\bar{Q}_S^2 = \frac{C_F}{N_c} Q_S^2 \quad (37)$$

and

$$Q_S^2 = \frac{8\pi^2\alpha N_c}{N_c^2 - 1} \sqrt{R^2 - b^2} \rho xG, \quad (38)$$

where $\bar{Q}_S$ is the quark saturation momentum and Q_S is the gluon saturation momentum. In (38) xG is the gluon distribution in a nucleon. Eq.(36) is a complete solution to the sea quark distribution of a nucleus in the one-quark-loop approximation.

Problem 9(M-H): Use (30) and (36) to show that[6]

$$\frac{dx(q_f + \bar{q}_f)_A}{d^2b d^2\ell} = \frac{N_c}{2\pi^4} \quad (39)$$

when $\ell^2/\bar{Q}_S^2 \ll 1$.

Eq.(39) gives meaning to the idea of saturation as a maximum occupation number for, in this case, quarks. As the nucleus gets larger and larger, that is as R grows, $\bar{Q}_S^2$ grows and so there are more and more sea quarks in the nuclear wavefunction, nevertheless, the 2-dimensional occupation number hits a constant upper bound for momenta below the saturation momentum.

Problem 10(M-H): Show that 1/2 of (39) comes from the first term in [] in (36) and that 1/2 comes from the second term with the third and

fourth term being small. The second term can be viewed as due to inelastic reactions while the first term is the elastic shadow of these inelastic reactions of a dipole passing over the nucleus.

9 Gluon Saturation in a Large Nucleus; the McLerran Venugopalan Model¹²

In order to directly probe gluon densities it is useful to introduce the “current”

$$j = -\frac{1}{4}F_{\mu\nu}^a F_{\mu\nu}^a. \quad (40)$$

We shall then calculate the process $j + A \rightarrow \text{gluon } (\ell) + \text{anything}$. I shall interpret the process in a slightly modified Bjorken frame, and in light-cone gauge, while we shall do the calculation in a covariant gauge and in the rest system of the nucleus[2].

For the interpretation we take the momenta of a nucleon in the nucleus and the current to be

$$p = (p + \frac{M^2}{2p}, 0, 0, p)$$

$$q = (0, 0, 0, -2xp).$$

For small x p is large since $p = \frac{Q}{2x}$ and this frame is much like an infinite momentum frame. If we choose an appropriate light-cone gauge, one that eliminates final state interactions, then the transverse momentum and the x distribution of gluons in the nuclear wavefunction is the same as the distribution of produced gluon jets labeled by ℓ in Fig.12.

In order to do the calculation of the spectrum of produced gluon jets we carry out a multiple scattering calculation, a term of which is illustrated in Fig.13[2]. The calculation is simplest to do in covariant gauge. Then

$$\frac{dx G_A}{d^2 b d^2 \ell} = \int_0^{2\sqrt{R^2 - b^2}} dz \rho x G(x, 1/\underline{x}^2) e^{-\frac{z}{L} \underline{x}^2 Q_s^2/4} e^{-i\ell \cdot \underline{x}} \frac{d^2 x}{4\pi^2} \quad (41)$$




FIG.12




FIG.13

where xG is the gluon distribution for a single nucleon, and we make use of the coordinate space interpretation where it corresponds to the gluon in the complex conjugate amplitude being separated from that in the amplitude by $\Delta x_{\perp} \sim 1/Q$. We note that the momentum space distribution of gluon produced off a single nucleon, the unintegrated gluon distribution, is given by

$$\frac{dxG}{d^2\ell} = \int \frac{d^2x}{4\pi^2} xG(x, 1/\underline{x}^2) e^{-i\ell \cdot \underline{x}} \quad (42)$$

as can be easily verified by integrating the lefthand side of (42) over $d^2\ell \Theta(Q^2 - \ell^2)$. Carrying out the z -integration in (4) one gets[13]

$$\frac{dxG_A}{d^2bd^2\ell} = \frac{N_c^2 - 1}{4\pi^4\alpha N_c} \int d^2x \frac{1 - e^{-\underline{x}^2 Q_S^2/4}}{\underline{x}^2} e^{-i\underline{\ell}\cdot\underline{x}} \quad (43)$$

where Q_S^2 is given in (38). We also note that the $e^{-\frac{z}{L}\underline{x}^2 Q_S^2/4}$ factor in (41) is just $S^\dagger(\underline{b})S(\underline{b}+\underline{x}) = S(\underline{x})$ for a gluon dipole to pass over a length z of nuclear material. Thus the derivation of (41) closely resembles that leading to (35) but now for gluons rather than quarks. Finally, when $\underline{\ell}^2/Q_S^2 \ll 1$ we find from (43)

$$\frac{dxG_A}{d^2bd^2\ell} \xrightarrow{\underline{\ell}^2/Q_S^2 \ll 1} \frac{N_c^2 - 1}{4\pi^3\alpha N_c} \ell n(Q_S^2/\underline{\ell}^2) \quad (44)$$

while for $\underline{\ell}^2/Q_S^2 \gg 1$ Eq.(43) gives the nuclear gluon distribution as simply a factor A times the nucleon gluon distribution. Eq.(44) shows that saturation is somewhat more complicated for gluons. The factor of $N_c^2 - 1$ counting the number of species of gluons is expected as is the factor of αN_c in the denominator. What is a little surprising is the log factor for which there is not yet a good intuitive understanding. Whether this log is a general factor or a peculiarity of the present model is not known for sure[6, 14].

10 The Golec-Biernat Wüsthoff Model¹⁵

We turn for a while to some phenomenology to see whether there is evidence for saturation of parton densities in deep inelastic lepton-proton scattering. So far the best way that has been found to approach this problem is through a simple model of deep inelastic and diffractive scattering inspired by the idea of saturation. We can motivate this discussion by going back to (36) and, supposing that such a picture might apply to a proton as well as a large nucleus, summing over f and integrating over $\underline{\ell}$ and $\underline{b}$ obtain

$$F_2(x, Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}} \int d^2x \int_0^1 dz \sum_\lambda |\psi_{T\lambda}^f(\underline{x}, z, Q)|^2 \int d^2b [1 - e^{-\underline{x}^2 \bar{Q}_S^2/4}]. \quad (45)$$

Now our $\bar{Q}_S^2$ naturally depends on the impact parameter b , as indicated in (37) and (38) for a nuclear target, however, as an approximation we suppose

$$\int d^2b[1 - e^{-\underline{x}^2 \bar{Q}_S^2/4}] = \sigma_0(1 - e^{-\underline{x}^2/4R_0^2}) \quad (46)$$

where R_0 will now be taken to depend only on x . Thus

$$F_2(x, Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}} \int d^2x \int_0^1 dz \sum_{\lambda} |\psi_{T\lambda}^f(\underline{x}, z, Q)|^2 (1 - e^{-\underline{x}^2/4R_0^2}) \sigma_0 \quad (47)$$

which is the formula used by Golec-Biernat and Wüsthoff[15]. In addition it is then natural to take the diffractive cross section to be given by the shadow of the inelastic collisions in which case one replaces $\sigma_0(1 - e^{-\underline{x}^2/4R_0^2}) = \sigma_0(1 - S)$ by $\frac{1}{2}\sigma_0(1 - S)^2$ giving

$$F_2^D(x, Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}} \int d^2x \int_0^1 dz \sum_{\lambda} |\psi_{T\lambda}^f(\underline{x}, z, Q)|^2 \frac{1}{2}\sigma_0(1 - e^{-\underline{x}^2/4R_0^2})^2. \quad (48)$$

Eq.(47) represents a total cross section, (48) represents an “elastic” cross section while the inelastic contribution would have a factor of $\frac{1}{2}\sigma_0(1 - e^{-\underline{x}^2/2R_0^2})$ replacing the last factors in (47) and (48).

Golec-Biernat and Wüsthoff include a quark-antiquark-gluon scattering term in addition to the quark-antiquark dipole term given by (48) so that larger mass diffractive states can also be described. For our purposes this is a detail which in any case introduces no new parameters. The model has three parameters

$$\sigma_0 = 23mb, \quad R_0^{-2} = \langle \bar{Q}_S^2 \rangle = \left(\frac{x_0}{x}\right)^\lambda GeV^2 \quad (49)$$

where $\lambda = 0.3$ and $x_0 = 3 \times 10^{-4}$. With these three parameters a good fit to low and moderate Q^2 and low x F_2 and F_2^D data is obtained. In fact the fit is surprisingly good over a range of Q^2 which is remarkably large given that there is no QCD evolution present in the model. We shall have to wait for further tests and refinements to be sure that the fits are meaningful, but we may have the first bit of evidence for saturation effects. The fact that $\langle \bar{Q}_S^2 \rangle$ is in the $1GeV^2$ region is reasonable.

11 Measuring Dipole Cross Sections

Refer back to (47). It is easy to check that

$$F_2 = c \int_{1/Q^2}^{1/Q_S^2} \frac{d\underline{x}^2}{\underline{x}^2} \quad (50)$$

when $Q^2/Q_S^2 \gg 1$. Thus, although we may view F_2 as being given by a dipole cross section the size of the dipole is not well determined by Q^2 but, rather, varies between $1/Q$ and $1/Q_S$. Thus currently deep inelastic structure functions are not well suited for determining dipole cross sections. The situation should improve considerably when the longitudinal structure function is measured, for in this case the dipole size will be fixed to be of size $1/Q$ and will give a direct measure of the dipole cross section.

Problem 11(E): Use 30 and (45) to derive (50). What is c ?

At present the best place to measure dipole cross sections, and thus to see how close or how far one is from finding unitarity limits, appears to be in the production of longitudinally polarized vector mesons[16].

The cross section for $\gamma_L^* + \text{proton} \rightarrow \rho_L + \text{proton}$ can be written as

$$\frac{d\sigma^{\gamma_L^* \rightarrow \rho}}{dt} = \frac{1}{4\pi} \left| \int d^2x d^2b \int_0^1 dz \psi_\rho^*(\underline{x}, z) (1 - S(\underline{x}, \underline{b})) \psi_{\gamma^*}(\underline{x}, z, Q)^{i\mathbf{b} \cdot \underline{\Delta}} \right|^2 \quad (51)$$

where $t = -\underline{\Delta}^2$ is the momentum transfer. The process is pictured in Fig.14 and can be viewed in three steps. (i) The virtual photon breaks up into a quark-antiquark dipole of size $\underline{x}$ with z being the longitudinal momentum fraction of the γ^* carried by either the quark or antiquark. (ii) The dipole scatters elastically on the proton with scattering amplitude $1 - S(\underline{x}, \underline{b})$ where $\underline{x}$ is the dipole size and $\underline{b}$ the impact parameter of the scattering. (iii) The quark-antiquark pair then become a ρ long after they have passed the proton. This is the sequence of steps given in (51) where one also integrates over all possible dipole sizes and where the integration over impact parameters gives a definite transverse momentum.

Now suppose S is purely real, and define $N(Q) = (\psi_\rho, \psi_{\gamma^*})$. Then one can take the square root of both sides of (51), and after taking the inverse Fourier transform one finds




FIG.14

$$\langle S(x, r_0, b) \rangle = 1 - \frac{1}{2N\pi^{3/2}} \int d^2\Delta e^{-i\vec{b}\cdot\vec{\Delta}} \sqrt{\frac{d\sigma}{dt}}. \quad (52)$$

In (52) $\langle S \rangle = (\psi_\rho, S\psi_{\gamma^*})$ while x denotes the Bjorken- x value of the scattering and $r_0(Q)$ denotes the typical value of the dipole size contributing to (51). For (52) to have meaning it is important that the range of dipole sizes contributing to (51) not be too large, and this appears to be the case for longitudinal ρ production. This analysis is modeled on the classic analysis of Amaldi and Shubert[17] for proton-proton elastic scattering. Thus one can estimate the S -matrix for a dipole of size r_0 , determined by Q , scattering on a proton at a given impact parameter if the data are good enough to carry out the integral in (52). The analysis is not model independent as one needs to take a wavefunction for the ρ [18, 19, 20] in order to evaluate N . However, N does not appear to be very sensitive to the choice of wavefunction. Also the data are not good enough at larger values of t to accurately determine S below $b \approx 0.3 fm$. One finds, for example, at $Q^2 = 3.5 GeV$ where $r_0 \approx 1/5 fm$ and for $x \approx 10^{-3}$ that: (i) $S(b \approx 0) \approx 0.5 - 0.7$; (ii) the probability of an inelastic collision $= 1 - S^2(b)$ is considerable at small values of b indicating a reasonable amount of blackness at central impact parameters; (iii) $\sigma_{tot}^{q\bar{q}-Proton} \approx 10 mb$; (iv) $\bar{Q}_S$ is consistent with that found in the Golec-Biernat Wüsthoff model. This adds, perhaps, another piece of evidence that saturation is approached in the HERA regime for moderate values of Q^2 .

12 A Simple Equation for Unitarity; the Kovchegov equation²¹

Consider a (not too high momentum) dipole scattering on a high-energy hadron. We suppose the quark-antiquark dipole is left moving while the hadron is right moving. Further we suppose that the rapidity, y , of the dipole is such that $\alpha y \ll 1$ so that one need not consider radiative corrections to the wavefunction of the dipole to evaluate the scattering amplitude. We wish to study the dependence of the elastic scattering amplitude as one changes the relative rapidity of the dipole and the hadron by an amount dY when the relative rapidity is Y . Clearly one can view the change dY either as increasing the momentum of the hadron and thus allowing its wavefunction to evolve further or as increasing the momentum of the dipole. The latter is easier to deal with since the dipole is a simple object. When the rapidity of the dipole is increased there is a small probability, proportional to dY , that the dipole emits a gluon before it scatters off the hadron. We now calculate the probability for producing this quark-antiquark-gluon state.

Since a gluon is emitted either off the quark or off the antiquark we have already done the basic emission amplitude, and it is given in (8). It will be convenient to work in a basis where transverse coordinate are used rather than transverse momenta so one must take the Fourier transform of (8). Thus the amplitude for a quark having transverse coordinate $\underline{x}_1$ to emit a gluon having transverse coordinate $\underline{z}$, longitudinal momentum k_- (Recall that in Sec.2 we were dealing with right movers while here we are concerned with left movers.) and polarization λ is

$$\psi_\lambda^c(\underline{z} - \underline{x}_1) = \int \frac{d^2k}{(2\pi)^2} e^{i(\underline{z} - \underline{x}) \cdot \underline{k}} \psi_\lambda^c(k) \quad (53)$$

with $\psi_\lambda^c(k)$ given by (8) with the replacement $k_+ \rightarrow k_-$. We are now using $A_- = 0$ light-cone gauge to evaluate the left moving quark-antiquark-gluon state. Using

$$\int \frac{d^2k}{2\pi} e^{i\underline{x} \cdot \underline{k}} \frac{\underline{k}}{k^2} = i \frac{\underline{x}}{\underline{x}^2} \quad (54)$$

one gets




FIG.15

$$\psi_\lambda^c(\underline{z} - \underline{x}_1) = \left(\frac{\lambda^c}{2}\right) 2ig \frac{(\underline{z} - \underline{x}_1) \cdot \underline{\epsilon}^{\lambda*}}{(\underline{z} - \underline{x}_1)^2} \frac{1}{\sqrt{(2\pi)^3 2k_-}}. \quad (55)$$

To calculate the probability of a quark-antiquark-gluon state one adds the graphs in Fig.15 to get[22]

$$dP_r = \sum_c \left(\frac{\lambda^c}{2} \frac{\lambda^2}{2}\right) 4g^2 \frac{d^2 z dk_-}{(2\pi)^3 2k_-} \left[-2 \frac{(\underline{z} - \underline{x}_1) \cdot (\underline{z} - \underline{x}_2)}{(\underline{z} - \underline{x}_1)^2 (\underline{z} - \underline{x}_2)^2} + \frac{1}{(\underline{z} - \underline{x}_1)^2} + \frac{1}{(\underline{z} - \underline{x}_2)^2} \right] \quad (56)$$

or

$$dP_r = \frac{\alpha N_c}{2\pi^2} d^2 z dY \frac{(\underline{x}_1 - \underline{x}_2)^2}{(\underline{x}_1 - \underline{z})^2 (\underline{x}_2 - \underline{z})^2} \quad (57)$$

where we have set $dk_-/k_- = dY$ and $C_F = \frac{N_c}{2}$ in the large N_c limit where the Kovchegov equation will be valid.

Then the S -matrix for the quark-antiquark-gluon state to elastically scatter on the hadron multiplied by dP_r in (57) gives the change in the S -matrix, dS , for dipole-hadron scattering. The result is the Kovchegov equation[21]

$$\frac{dS(\underline{x}_1 - \underline{x}_2, Y)}{dY} = \frac{\alpha N_c}{2\pi^2} \int d^2 z \frac{(\underline{x}_1 - \underline{x}_2)^2}{(\underline{x}_1 - \underline{z})^2 (\underline{x}_2 - \underline{z})^2} [S(\underline{x}_1 - \underline{z}, Y) S(\underline{z} - \underline{x}_2, Y) - S(\underline{x}_1 - \underline{x}_2, Y)], \quad (58)$$

$$\frac{DS}{dY} = \begin{array}{c} \text{---} \overline{x_1} \text{---} \\ \text{---} \overline{z} \text{---} \\ \text{---} x_2 \text{---} \end{array} + \begin{array}{c} \text{---} \overline{x_1} \text{---} \\ \text{---} \overline{z} \text{---} \\ \text{---} x_2 \text{---} \end{array} + \begin{array}{c} \text{---} \overline{x_1} \text{---} \\ \text{---} \overline{z} \text{---} \\ \text{---} x_2 \text{---} \end{array}$$

FIG.16

and it is illustrated in Fig.16. We have assumed that the scattering of the two dipoles, the quark-(antiquark part of the gluon) and the (quark part of the gluon)-antiquark dipoles, factorize when scattering off the hadron. This was clear in the model Kovchegov considered where the hadron was a large nucleus. This factorization is less obvious in the general case and the Kovchegov equation may be a sort of mean field approximation to a more complete equation. Also the final term on the right-hand side of (58), corresponding to the last two graphs in Fig.16, give the virtual contributions necessary to normalize the wavefunction[22]. The necessity of this last term can be seen by considering the weak interaction limit where $S \rightarrow 1$. Then the final term on the righthand side of (58) is necessary to get $\frac{dS}{dY} = 0$ when $S = 1$.

There are two interesting limits to (58). First suppose that S is near 1 and write $S = 1 - iT$. One easily finds, keeping only linear terms in T ,

$$\frac{dT(\underline{x}_1 - \underline{x}_2, Y)}{dy} = \frac{\alpha N_c}{\pi^2} \int d^2 z \frac{(\underline{x}_1 - \underline{x}_2)^2}{(\underline{x}_1 - \underline{z})^2 (\underline{x}_2 - \underline{z})^2} [T(\underline{x}_1 - \underline{z}, Y) - \frac{1}{2} T(\underline{x}_1 - \underline{x}_2, Y)] \quad (59)$$

which is the dipole form of the BFKL equation. In this case the factorized form of the scattering is justified by the large N_c limit and the weak coupling approximation.

The other interesting limit is where S is small in which case one need only keep the second term on the righthand side of (58) giving

$$\frac{dS(\underline{x}_1 - \underline{x}_2, Y)}{dY} = -\frac{\alpha N_c}{2\pi^2} \int \frac{d^2 z (\underline{x}_1 - \underline{x}_2)^2}{(x_1 - z)^2 (x_2 - z)^2} S(\underline{x}_1 - \underline{x}_2, Y). \quad (60)$$

Of course (60) as written cannot be valid. The assumption that S be small can be true only when the dipole size is large compared to $1/Q_s$. Thus we should restrict the integration in (60) to the region $(\underline{x}_1 - \underline{x}_2)^2 \gg 1/Q_s^2$ as well as to the region $(\underline{x}_1 - \underline{z})^2, (\underline{x}_2 - \underline{z})^2 \gg 1/Q_s^2$ so that the nonlinear term in (58) not cancel the linear term. In the logarithmic regions of integration one can rewrite the integral (60) as

$$\frac{dS(\underline{x}_1 - \underline{x}_2, Y)}{dY} = -2 \frac{\alpha N_c}{2\pi^2} \int_{1/Q_s^2}^{(\underline{x}_1 - \underline{x}_2)^2} \frac{\pi d(\underline{x}_1 - \underline{z})^2}{(\underline{x}_1 - \underline{z})^2} S(\underline{x}_1 - \underline{x}_2, Y) \quad (61)$$

giving

$$\frac{dS(\underline{x}, Y)}{dY} = -\frac{\alpha N_c}{\pi} \ell n(Q_s^2 \underline{x}^2) S(\underline{x}, Y) \quad (62)$$

whose solution is

$$S(\underline{x}, Y) = e^{-\frac{\alpha N_c}{\pi} \int_{Y_0}^Y dy \ell n[Q_s^2(y) \underline{x}^2]} S(\underline{x}, Y_0). \quad (63)$$

If Q_s^2 is exponentially behaved

$$Q_s^2(y) = e^{c \frac{\alpha N_c}{\pi} (y - Y_0)} Q_s^2(Y_0)$$

then

$$S(\underline{x}, Y) = e^{-\frac{c}{2} (\frac{\alpha N_c}{\pi})^2 (Y - Y_0)^2} S(\underline{x}, Y_0) \quad (64)$$

where Y_0 should be chosen to satisfy

$$Q_s^2(Y_0) \underline{x}^2 = 1. \quad (65)$$

Eq.(64) is the result found in Ref.23 and argued heuristically already some time ago[24], although without an evaluation of the coefficient of the Y^2 term in the exponent.

The Kovchegov equation is an interesting equation for studying scattering when one is near unitarity limits. In the next section we shall use a procedure very similar to what we have done here to derive an equation whose content is presumably equivalent to the Balitsky equation[5], a somewhat more general form than the Kovchegov equation. The advantage of (58) is its simplicity and its likely qualitative correctness in QCD.

13 A Simple Derivation of the JIMWLK equation^[14,25–27]

Over the past seven years or so there has been an ambitious program dedicated to finding appropriate equations for dealing with high density wavefunctions in QCD. This program has been quite successful and a renormalization group equation in the form of a functional Fokker-Planck equation for the wavefunction of a high-energy hadron has been given by the authors of Refs.14 and 25-27. (JIMWLK). The most complete derivation is given in Ref.27 where the equation is written in terms of a covariant gauge potential, α , coming from light-cone gauge quanta in a high-energy hadron. Here we give an alternative simple derivation[28].

We can imitate this mixture of gauges used in Ref.27 by taking Coulomb gauge[29] which has a gluon propagator

$$D_{\alpha\beta}(k) = -\frac{i}{k^2} \left[g_{\alpha\beta} - \frac{N \cdot k (N_\alpha k_\beta + N_\beta k_\alpha) - k_\alpha k_\beta}{(\vec{k})^2} \right] \quad (66)$$

where $N \cdot v = v_0$ for any vector v . Suppose the propagator has $k_+^2 \gg \underline{k}^2, k_-^2$ and connects two highly right moving lines. Then the Coulomb gauge propagator is equivalent to the $A_+ = 0$ light-cone gauge propagator

$$D_{\alpha\beta}(k) = \frac{-i}{k^2} \left[g_{\alpha\beta} - \frac{\eta_\alpha k_\beta + \eta_\beta k_\alpha}{\eta \cdot k} \right] \quad (67)$$

Problem 12(E): Show that D_{--} , and D_{-i} as given in (66) and (67) agree when $\underline{k}_+^2 \gg \underline{k}^2, k_-^2$. For a right moving system these are the important components of the gluon propagator.

Similarly for a left moving system Coulomb gauge is equivalent to $A_- = 0$ gauge while for gluon lines which connect left moving systems to right moving systems the dominant component in (66) is $D_{+-} = 1/(\vec{k})^2$ which looks like covariant gauge when $k_+^2, k_-^2 \ll \underline{k}^2$.

We are going to consider the scattering of a set of left moving quanta, quarks and gluons, on some high-energy, right moving hadron. These quarks and gluons may be parts of a hadronic wavefunction which are frozen in the passage over the right moving hadron or they may come from a current as in our discussion of deep inelastic scattering given in Sec.8. For simplicity we shall limit our discussion to left moving quark and antiquark lines, but

this is simply to avoid too cumbersome notation. Then a left moving quark interacting with the right moving hadron can be represented by

$$V^\dagger(\underline{x}) = P \exp\{ig \int_{-\infty}^{\infty} dx_- A_+(\underline{x}, x_-) \} \quad (68)$$

where we have taken $x_+ = 0$ and fixed the left moving quark to have transverse coordinate $\underline{x}$. Except for the change of right moving quark to left moving quark (68) is the same as the operator in the matrix element in (26) where we showed how quarks could be identified with Wilson lines in the fundamental representation. By taking gauge invariant combinations of V 's and $V^\dagger$'s we can form observables which depend on A_+ and which correspond to the scattering of quite general left moving systems on the right moving hadron. We denote a general such observable by $O(A_+)$.

Although we have put the integration in (68) exactly on the light-cone we in fact are going to assume that the left moving observable has rapidity y obeying $\alpha y \ll 1$ so that transverse gluons are unlikely to be emitted by the left movers allowing us to identify the left moving system at $x_- = \pm\infty$. If the relative rapidity of the scattering is Y then we imagine that $\alpha(Y - y) \approx \alpha Y \gg 1$ so that the right moving hadron has, in general, a wavefunction including many gluons. If the right moving hadron has momentum p then the scattering amplitude is

$$\langle O \rangle_Y = \langle p | O | p \rangle = \int D[\alpha(\underline{x}, x_-)] O(\alpha) W_Y[\alpha] \quad (69)$$

where the weight function W_Y is given by

$$W_Y[\alpha] = \int D[A_\mu] \delta(A_+ - \alpha) \delta(F(A)) \Delta_F[A] e^{iS[A]} \quad (70)$$

where D indicates a functional integral, F is a gauge fixing, and Δ_F is the corresponding Fadeev-Popov determinant times an operator which projects out the state $|p\rangle$ initially and finally. We suppose W_Y is normalized to

$$\int D[\alpha] W_Y[\alpha] = 1. \quad (71)$$

Now consider the Y -dependence of $\langle O \rangle_Y$. From (69) one can clearly write

$$\frac{d}{dY} \langle O \rangle_Y = \int D[\alpha] O(\alpha) \frac{d}{dY} W_Y[\alpha]. \quad (72)$$




FIG.17

However, one can equally well imagine calculating $\frac{d}{dY} \langle O \rangle_Y$ by evaluating the change of the left moving system, due to an additional gluon emission, as one increases the rapidity of the left moving system by an amount dY . The change in the left moving state is given by the gluon emissions and absorptions shown in Figs.17 and 18 where spectator quark and antiquark lines are not shown. In Fig.17 we have assumed that a gluon connects a quark and an antiquark line. This is for definiteness. We equally well could have assumed a connection to two quark lines or to two antiquark lines. The vertical line in the figures represents the “time,” $x_- = 0$, at which the left moving system passes the right moving hadron. This view of the Y -dependence of $\langle O \rangle_Y$ in terms of a change of rapidity of the (rather simple) left movers is in the spirit of work previously done by Balitsky[5], Kovchegov[21], and Weigert[26].

We shall examine in some detail the graphs of Fig.17 before stating the complete result including the graphs of Fig.18. Begin with the graph shown in Fig.17c. We do the calculation in $A_- = 0$ light-cone gauge, which for left movers is equivalent to our Coulomb gauge choice. This graph is exactly the same as has been calculated in Sec.11 except for the fact that the quark and antiquark lines are not necessarily in a color singlet. The result is the same as $\frac{1}{2}$ the first term on the righthand side of (56) except for the color factors which we put in separately. The result is




FIG.18

$$V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2) \rightarrow -\frac{\alpha_S}{\pi^2} \int d^2 z \frac{(\underline{x}_1 - \underline{z}) \cdot (\underline{x}_2 - \underline{z})}{(\underline{x}_1 - \underline{z})^2 (\underline{x}_2 - \underline{z})^2} \tilde{V}_{cd}(\underline{z}) V^\dagger(\underline{x}_1) T^c \otimes V(\underline{x}_2) T^d \quad (73)$$

in going from O to $\frac{dO}{dY}$. The additional factors, as compared to the corresponding term in (56), are T^c which comes to the right of $V^\dagger(\underline{x}_1)$ because the emission off the $\underline{x}_1$ -line is at early values of x_- , the T^d which comes to the right of $V(\underline{x}_2)$ because the absorption on the $\underline{x}_2$ -line is at late values of x_- , and the factor $\tilde{V}_{cd}(\underline{z})$ giving the interaction of the gluon with the hadron as a Wilson line in the adjoint representation.

Now write

$$V^\dagger(\underline{x}_1) T^c = V^\dagger(\underline{x}_1) T^c V(\underline{x}_1) V^\dagger(\underline{x}_1) = \tilde{V}_{ca}(\underline{x}_1) T^a V^\dagger(\underline{x}_1). \quad (74)$$

Now T^a comes to the left of $V^\dagger(\underline{x}_1)$ as if the emission of the gluon were at late values of x_- . Indeed, we may view the graph in Fig.17c as being given by the “mnemonic” graph shown in Fig.19 where the adjoint line integral starts at large positive values of x_- and proceeds to large negative values of x_- at a transverse coordinate $\underline{x}_1$ then back again to large positive values of x_- at a transverse coordinate $\underline{z}$. The result of the graph of Fig.17c then is

$$V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2) \rightarrow \frac{\alpha_S}{\pi^2} \int d^2 z \frac{(\underline{x}_1 - \underline{z}) \cdot (\underline{x}_2 - \underline{z})}{(\underline{x}_1 - \underline{z})^2 (\underline{x}_2 - \underline{z})^2} \{-\tilde{V}^\dagger(\underline{x}_1) \tilde{V}(\underline{z})\}_{ab} T^a V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2) T^b \quad (75)$$

Now it is straightforward to add in the other terms of Fig.17 and those of Fig.18 to get




FIG.19

$$V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2) \rightarrow \alpha_S \left\{ \frac{1}{2} \int d^2x d^2y \eta_{\underline{x}\underline{y}}^{ab} \frac{\delta^2}{\delta\alpha^a(\underline{x}, x_-) \delta\alpha^b(\underline{y}, y_-)} + \int d^2x \nu_{\underline{x}}^a \frac{\delta}{\delta\alpha^a(\underline{x}, x_-)} \right\} \cdot V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2) \quad (76)$$

where, if one takes large positive values of x_- and y_- , the functional derivatives in (76) simply insert color matrices in the appropriate places, as, for example, in (75). η and ν are given by

$$g^2 \eta_{\underline{x}\underline{y}}^{ab} = 4 \int \frac{d^2z}{4\pi^2} \frac{(\underline{x} - \underline{z}) \cdot (\underline{y} - \underline{z})}{(\underline{x} - \underline{z})^2 (\underline{y} - \underline{z})^2} \cdot \{1 + \tilde{V}^\dagger(\underline{x}) \tilde{V}(\underline{y}) - \tilde{V}^\dagger(\underline{x}) \tilde{V}(\underline{z}) - \tilde{V}^\dagger(\underline{z}) \tilde{V}(\underline{y})\}_{ab} \quad (77)$$

and

$$g\nu_{\underline{x}}^a = 2i \int \frac{d^2z}{(\underline{x} - \underline{z})^2} \text{tr}[T^a \tilde{V}^\dagger(\underline{x}_1) \tilde{V}(\underline{z})]. \quad (78)$$

For a general scattering one simply replaces $V^\dagger(\underline{x}_1) \otimes V(\underline{x}_2)$ in (76) by $O(\alpha)$. After multiplying by $W_Y[\alpha]$ and integrating over $D[\alpha]$ one gets[28]

$$\int D[\alpha] O(\alpha) \frac{dW_Y[\alpha]}{dY} = \int D[\alpha] O(\alpha) \alpha_S \left\{ \frac{1}{2} d^2x d^2y \frac{\delta^2}{\delta\alpha^a(\underline{x}, x_-) \delta\alpha^b(\underline{y}, y_-)} \cdot [W_Y \eta_{\underline{x}\underline{y}}^{ab}] - \int d^2x \frac{\delta}{\delta\alpha^a(\underline{x}, x_-)} [W_Y \nu_{\underline{x}}^a] \right\} \quad (79)$$

where an integration by parts in α has been done on the righthand side of (79). To the extent that the $O(\alpha)$ form a complete set of observables, and it is not clear how close this is to being true, one can equate the integrands of (79) and arrive at the JIMWLK equation

$$\begin{aligned} \frac{dW_Y[\alpha]}{dY} = \alpha_S \{ & \frac{1}{2} \int d^2x d^2y \frac{\delta^2}{\delta\alpha^a(\underline{x}, x_-) \delta\alpha^b(\underline{y}, y_-)} [W_Y \eta_{\underline{x}\underline{y}}^{ab}] \\ & - \int d^2x \frac{\delta}{\delta\alpha^a(\underline{x}, x_-)} [W_Y \nu_{\underline{x}}^a] \}. \end{aligned} \quad (80)$$

The exact values of x_- and y_- appear to have some arbitrariness as discussed in some detail in Ref.28. Eq.(80) is an elegant equation of a functional Fokker-Planck type the nature of whose solutions is now under investigation.

References

- [1] Yu.V. Kovchegov, *Phys.Rev.D***54** (1996) 5463; **D55** (1997) 5445.
- [2] Yu. V. Kovchegov and A.H. Mueller, *Nucl. Phys.***B529** (1998) 451.
- [3] W. Buchmüller, M.F. McDermott and A. Hebecker, *Nucl.Phys.* **B487** (1997) 283; **B500** (1997) 621 (E)
- [4] W. Buchmüller, T. Gehrman and A. Hebecker, *Nucl.Phys.* **B538** (1999) 477.
- [5] I. Balitsky, *Nucl. Phys.***B463** (1996) 99.
- [6] A.H. Mueller, *Nucl.Phys.* **B558** (1999) 285.
- [7] J.D. Bjorken in *Proceedings of the International Symposium on Electron and Photon Interactions at High Energies*, pages 281-297, Cornell (1971).
- [8] A.H. Mueller, *Nucl. Phys.***B335** (1990) 115.
- [9] L.L. Frankfurt and M. Strikman, *Phys.Rep.***160** (1998) 235.
- [10] B. Blättel, G. Baym, L.L. Frankfurt and M. Strikman, *Phys. Rev.Lett.* **70** (1993) 896.

- [11] B.G. Zakharov, *JETP Lett.***63** (1996) 952.
- [12] L. McLerran and R. Venugopalan, *Phys. Rev.***D49**(1994) 2233; **D49**(1994) 3352; **D50**(1994) 2225.
- [13] J. Jalilian-Marian, A. Kovner, L. McLerran and H. Weigert, *Phys. Rev.* **D55** (1997) 5414.
- [14] E. Iancu and L. McLerran, *Phys. Lett.***B510** (2001) 145.
- [15] K. Golec-Biernat and M. Wüsthoff, *Phys. Rev.***D59** (1999) 014017; *Phys. Rev.***D60** (1999) 114023.
- [16] S. Munier, A.M. Stasto and A.H. Mueller, *Nucl. Phys.* **B603** (2001) 427.
- [17] U. Amaldi and K. R. Schubert, *Nucl. Phys.***B166** (1980) 301.
- [18] H.G. Dosch, T. Gousset, G. Kulzinger and H.J. Pirner, *Phys. Rev.***D55** (1997) 2602.
- [19] J. Nemchik, N.N. Nikolaev, E. Predazzi and B.G. Zakharov, *Z. Phys.***C75** (1997) 71.
- [20] L. Frankfurt, N. Koepf and M. Strikman, *Phys. Rev.***D54** (1996) 3194.
- [21] Yu. Kovchegov, *Phys. Rev.***D60** (1999) 034008; *Phys. Rev.***D61** (2000) 074018.
- [22] A.H. Mueller, *Nucl. Phys.* **B415** (1994) 373.
- [23] E. Levin and K. Tuchin, *Nucl. Phys.***B573** (2000) 83; hep-ph/0101275.
- [24] A.H. Mueller and G.P. Salam, *Nucl. Phys.***B475** (1996) 293.
- [25] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *Nucl. Phys.***B504** (1997) 415; *Phys. Rev.* **D59** (1999) 014014.
- [26] H. Weigert, hep-ph/0004044.
- [27] E. Iancu, A. Leonidov and L. McLerran, *Nucl. Phys.* **A692** (2001) 583; *Phys. Lett.***B510** (2001) 133; hep-ph/0109115.

- [28] A.H. Mueller, hep-ph/0110154.
- [29] T. Jaroszewicz, *Acta Physica Polonica* **B11** (1980) 965.